

HIV/STI Prevention and Young Men Who Have Sex with Men

The vital importance of reaching the nation's most at-risk population

Men who have sex with men (MSM) accounted for 65 percent of all new infections in the United States in 2013. Among young men ages 13-24, 90 percent of HIV infections were attributed to male-to-male sexual contact.¹ Young men who have sex with men (YMSM) are the only risk group in which the number of new infections has increased steadily each year since the 1990s, even as it has decreased among other populations.² Thirty years into the HIV epidemic, young men who have sex with men remain at serious risk of acquiring HIV.

Stigma and historical oppression complicate the problem, placing young men of color who have sex with men at even higher risk: the rate of new infections among this group has risen steadily for 20 years, including a 22 percent rise in new infections between 2008-2010. Black young people are also at high-risk, accounting for 67 percent of new HIV cases among young people, while young Latino men account for 17 percent of new HIV cases among young people. Each day, nine young Black men are diagnosed with HIV.

Without culturally-appropriate, pragmatic, and inclusive prevention strategies, YMSM are left to explore the realm of sex and sexuality uninformed and through trial and error – leaving them at risk.³ This document explores the barriers to HIV prevention YMSM may face and provides guidelines for creating more effective prevention programs.

STRUCTURAL BARRIERS PUT YMSM OF COLOR AT HIGHER RISK

- For many YMSM, social and economic factors, including homophobia, stigma, and lack of access to culturally competent health care and health care services may increase risk behaviors or be a barrier to receiving HIV prevention services.⁴
- According to the Centers for Disease Control and Prevention (CDC), 58 percent of all HIV/AIDS diagnoses among YMSM aged 13-24 in 2011 were among Black youth, even though Blacks represented only 17 percent of the population in that age group.⁵

- Studies have shown that among men who have sex with men, Black men are more at risk for HIV even when they have the same or fewer risk behaviors. An analysis of 53 studies found that Black men were not more likely than whites to have unprotected anal sex, engage in commercial sex work, or have sex with a known HIV positive partner; and in fact reported having fewer partners than white men – yet acquire HIV at vastly disproportionate rates. Their heightened risk for HIV is not due to more risk behaviors but is because the communities from which they are likely to select romantic and sexual partners experience higher rates, making their odds of acquiring the disease higher.⁶

- A history of oppression of Blacks dating back to slavery, and inequity that continues to the present impact HIV risk for Black youth. Systematic discrimination contributes to and is intertwined with underemployment and unemployment, decreased access to medical care, and incarceration; these along with social stigma around being LGBTQ create a strong barrier to sexual health for Black YMSM.¹⁴ For instance:

- HIV positive Black men who have sex with men are less likely than HIV positive white men to be taking ART (anti-retroviral therapy, a group of HIV medications). Because treatment for HIV decreases the viral load, those who are not receiving treatment are more likely to transmit HIV.¹⁶

- Black men are nine times as likely to be incarcerated as whites; incarcerated men are at risk for unprotected sex and HIV transmission.¹⁴

- More than one-quarter of Blacks live in poverty compared to 11 percent of whites.⁷ Blacks are less likely than whites to be insured and more likely to have publicly funded insurance.⁸

- Among YMSM ages 13-24, Latinos experienced 20 percent of new HIV infections in 2010. One study found that

FROM RESEARCH
TO PRACTICE

“Without culturally-appropriate, pragmatic, and inclusive prevention strategies, YMSM are left to explore the realm of sex and sexuality uninformed and through trial and error – leaving them at risk.”³”

cultural expectations play a large role in HIV risk for this population: men who held strong “machismo” (traditionally masculine gender-role) beliefs were more likely to have had multiple partners, while men who had experienced discrimination based on their sexual orientation were more likely to have had unprotected anal intercourse (UAI).⁹ Latinos are more than twice as likely to have no health insurance as non-Latinos, decreasing their access to health care.¹⁰

- American Indians and Alaska Natives represent less than one percent of the total number of HIV/AIDS cases reported to CDC’s HIV/AIDS Reporting System, but their rate of diagnosis of HIV is higher than that of whites (11.9 cases per 100,000 compared to 8.2).² Among males, male to male sexual contact accounted for 75 percent of cumulative HIV cases among American Indians and Alaska Natives in 2011 (the most recent year for which this data is available).¹¹ Many factors, including lack of reporting and cultural barriers to open discussion of same-gender sexual behavior, make understanding the HIV epidemic among this population a challenge.

SILENCE, STIGMA, AND DISCRIMINATION CONTRIBUTE TO HIV RISK AMONG YMSM

- YMSM not only face a lack of information about safer sex, but face strong barriers to developing self-efficacy for safer sex behaviors. Stigma against being LGBTQ leads to a dearth of information and limited discussion about safer sex and HIV prevention.
- All YMSM experience homophobia, through laws and policies that discriminate against them and harmful cultural messages. Most YMSM students have also directly experienced discrimination in schools: 92 percent of LGBTQ students report being verbally harassed in school, and 45 percent report being physically harassed in school.¹²
- LGBTQ youth are overrepresented among homeless youth – comprising approximately 40 percent. These youth are more likely to participate in sex work and

drug use and have very limited access to health care, placing them at grave risk of HIV.¹³

- Many students encounter misinformation and/or harmful stereotypes in HIV prevention education and sexuality education. Abstinence-only programs often rely on stereotypical gender roles and heterosexual relationships as models, not only ignoring LGBTQ youth, but contributing to stigma against those YMSM who don’t fit a traditionally masculine ideal.¹⁴ Even otherwise comprehensive sexuality education may fail to provide instruction on risk reduction for LGBTQ youth.³
- Other sources of information about safer sex, like parents and health care providers, may not be available to LGBTQ young people. Their parents may not be accepting of their identity or may not be prepared to discuss safer sex for LGBTQ individuals; and many are concerned their health care provider will not maintain confidentiality.³
- HIV-related stigma and homophobia are closely linked due to public perception of HIV and AIDS as a “gay disease.”¹⁵ One study found that 59 percent of men who had never been tested for HIV had not done so out of fear of negative social consequences.¹⁶ Another found that internalized homophobia was linked to lack of awareness of HIV prevention programs and to a lack of comfort with condom use.¹⁷

MANY YMSM ENGAGE IN BEHAVIORS THAT PUT THEM AT RISK OF HIV/STIS

- Determining the risky behaviors in which YMSM are likely to engage can inform behavioral interventions in communities of at-risk MSM and secondary prevention efforts among those already living with HIV.
- According to the National HIV Behavioral Surveillance System, 89 percent of YMSM reported anal intercourse with a male partner in the past year and 54 percent had unprotected anal intercourse (UAI). Seventeen percent had UAI with more than one male partner.¹⁸
- Findings from the same study show that compared to young men who had UAI with only one male partner, those who had UAI with multiple male partners were more likely to have engaged in UAI with a casual male partner, as opposed to someone with whom they were in a relationship. Thirty-one percent of all young men participating in the National HIV Behavior Surveillance System reported drug use during sex.⁴
- One study of YMSM found that the odds of HIV infection increased significantly as the age of one’s sexual partners increased – participants with partners five or more years older than they had twice the odds, and those with partners 10 years or older were 4 times as likely of becoming infected with HIV than study participants as a whole.¹⁹

- Many YMSM are not aware of their HIV status. In one nationwide study of MSM which included HIV testing, 19 percent of YMSM tested positive, with 44 percent of those who tested positive unaware of being infected with HIV.²⁰
- Studies have found that because of the existence of medications which can prolong an HIV positive individual's life and prevent the virus from developing into AIDS, some YMSM may be taking more sexual risks, including unprotected sex with untested or known HIV positive individuals.²¹

EFFECTIVE PROGRAMS, WHICH BUILD SKILLS AND AFFIRM THE VALUE OF YMSM, CAN REDUCE SEXUAL RISK-TAKING

Complex issues are fueling HIV transmission among YMSM—particularly for YMSM of color. In order to address those complexities, interventions must address individual behavior and the socio-cultural determinants that fuel HIV transmission.

- Peer-based interventions have been effective with YMSM in reducing risk behaviors.
 - One intervention with gay men significantly reduced sexual risk-taking behavior in four cities by recruiting popular peers and training them to pass on behavior recommendations to friends through conversation. Surveys found that at one-year follow-up, unprotected anal intercourse in the cities decreased between 15-29 percent, condom use increased, and the number of sex partners decreased.²²
 - One study found that YMSM were most likely to be reached effectively through outreach activities, such as dances, movie nights, gay rap/hip hop groups, and volleyball.²³
 - Another program showed 60 percent fewer YMSM reporting unprotected anal intercourse after sustained sexuality-related peer education that combined education sessions with a social support group.²⁴
 - One effective program presented a “menu” of risk reduction options over the course of a retreat weekend, delivered by trained peer facilitators. Compared to those who did not receive the intervention, participants reduced their number of sex partners and instances of UAI, and were 81 percent more likely to have been tested for HIV six months after the program.²⁵
- Because of the barriers posed by homophobia and racism, interventions for YMSM of color may need to focus on community-building approaches that reflect cultural nuances as well as on individual behavior change. For example, one study suggests that interventions focus on increasing the collective capacity of Black YMSM to address HIV and on increasing acceptance of YMSM within Black communities.^{26,27}
- One study found that programs that promoted positive youth development are paramount to fostering healthy sexual outcomes for youth. The study found that programs that build skills, enhance bonding, strengthen families, engage youth, and empower youth were the most effective at creating behavior change.²⁸

The following critical components for HIV/STI prevention are drawn from research.

- **Tailor programs to include YMSM.** Programs developed for all young people should discuss sexuality and should include discussion about anal sex in HIV/STD risk reduction sessions.²⁹
- **Involve youth.** Peer support groups and Gay/Straight alliances provide non-sexual opportunities for YMSM to share their emotions and experiences, ease their feelings of isolation, and build support systems. Involving YMSM in program design and implementation reduces their risky behaviors and fosters their spirit of self-determination and self-worth.^{25,30}
- **Utilize peer influence within social networks.** Research has shown that identifying and recruiting peer leaders is an effective way of reaching YMSM with HIV prevention messages.^{24,25}
- **Explore new venues, including non-gay-identified ones.** Community gathering places, welcoming churches, barber shops and beauty shops, websites, and bars or coffeehouses should be considered as a means of reaching YMSM with prevention messages.¹¹
- **Foster a sense of personal worth.** Prevention must affirm the value of YMSM and create a context that fosters responsible sexual behavior. One-on-one counseling sessions make effective beginnings for such interventions.^{3,11,13}
- **Address the needs of youth.** Focus on the needs identified by YMSM, not on those perceived by adults. This may include sponsoring support groups, building dating skills, and providing mentors and other role models.²⁹
- **Teach skills.** Programs must teach skills. The ability to use condoms, negotiate safer sex with partners, build relationships, communicate with steady and casual partners, make decisions, and say ‘no’ strengthens teens in making healthy choices.²⁹

“Prevention must affirm the value of YMSM and create a context that fosters responsible sexual behavior.”

- **Incorporate risk reduction strategies.** Programs should include information about partner reduction, gauging one's own susceptibility, the relative risk of specific behaviors (e.g., anal sex vs. oral sex), and other ways to reduce HIV risk beyond abstinence and condom use. Recent studies show the promise of preexposure prophylaxis (PrEP) for preventing HIV infection, but research efforts suffer from disproportionately low representation of the youth who are most at risk. Youth-focused research is critical and should include behavioral, community, and biomedical interventions to create a comprehensive HIV prevention package.³²
 - **Provide sustained support.** Since sustaining behavior change is difficult, populations at high risk require continuing support and reinforcement. To prevent relapse into unsafe behavior, prevention programs must address the changing needs of YMSM as they grow older.
 - **Create programs specifically for YMSM of color.** Studies indicate that programs need to address individual, community, and cultural factors pertinent to YMSM. Programs should address racism in the gay, white community while simultaneously supporting YMSM of color as they deal with decisions regarding sexuality, gay identity, culture, and race/ethnicity. YMSM also need safe environments for sharing their experiences.²⁹
 - **Address the needs of marginalized groups such as homeless youth and IV drug users.** Programs must reach out to homeless youth, especially those involved in commercial sex, and those who are IV drug users. One study of homeless adolescents found that one in 14 had been treated for HIV.¹³ An estimated one-third of homeless youth have participated in survival sex or sex work. Homeless youth are also more vulnerable to intravenous drug use (IDU), a risk factor for HIV transmission. And in 2011, 11 percent of new HIV infections among males were attributed to IDU or male sexual contact and IDU. This highly at-risk population desperately needs outreach programs and basic health care.¹⁹
- Importantly, interventions which focus solely on changing personal behaviors are only a part of a successful HIV prevention strategy. Addressing other factors which contribute to the spread of HIV is vital. Recommendations drawn from research include:
- **Support structural interventions (approaches that promote health by changing the environment to one that facilitates health), including ones which work toward:**
 - Population education - educational techniques designed to raise the consciousness of its participants and allow them to become more aware of how an individual's personal experiences are connected to larger societal problems.
 - Harm reduction, including needle exchange legislation and programs for injection drug users.
 - Community mobilization for HIV prevention advocacy and a reduction of stigma against HIV positive individuals
 - Ensuring access to quality education and health care
 - Routine HIV testing
 - Engaging families in safer sex programs
 - Reducing the community viral load through early diagnosis of HIV infection, timely initiation of antiretroviral therapy, and treatment adherence.
 - Supporting policies protecting against hate crimes and employment discrimination based on sexual orientation.³²
 - **Adopt an ecological approach to prevention.** An ecological approach attempts to create more effective and culturally competent programs by examining a young person's entire sphere (their family, community, and societal relationships and influences) and creating peer, group and family-level interventions.³³

CONCLUSION

Young men who have sex with men are highly at risk for HIV. Like all young people, YMSM need culturally-appropriate, pragmatic, and inclusive prevention messages – as well as programs which address behavioral, cultural, and institutional barriers to sexual health information and services.

Written by Durryle Brooks, Program Manager, GLBTQ Initiatives, and Emily Bridges, Director of Public Information Services

Updated by Wesley Thomas, Program Coordinator for GLBTQ Health and Rights, 2015

Advocates for Youth © August 2015

REFERENCES

1. CDC. "Slide Show: HIV Surveillance in Men who Have Sex with men." Accessed from <http://www.cdc.gov/hiv/topics/surveillance/resources/slides/msm/index.htm> on August 6, 2014.
2. Centers for Disease Control and Prevention (CDC). Subpopulation Estimates From the HIV Incidence Surveillance System -- United States, 2006. MMWR. 2008; 57(36):985-989
3. Kubicek K. "In the Dark: Young Men's Stories of Sexual Initiation in the Absence of Relevant Sexual Health Information." Health Education Behavior April 2010; 37(2) 243-263.

4. Mechoulam, S. (2011). The external effects of black male incarceration on black females. *Journal of Labor Economics*, 29(1), 1-35.
5. U.S. Census Bureau. 2009 American Community Survey 1-Year Estimates, Detailed Tables. Accessed from factfinder.census.gov on October 26, 2010.
6. Millett et al. Explaining disparities in HIV infection among black and white men who have sex with men: a meta-analysis of HIV risk behaviors. *AIDS* 21 (15) 2083-2091.
7. U.S. Census Bureau. Detailed Tables: Poverty. Accessed from factfinder.census.gov on November 8, 2010.
8. Kaiser State Health Facts. "Distribution of the Nonelderly Uninsured by Race/Ethnicity, states (2008-2009), U.S. (2009)." Kaiser Family Foundation. Accessed from <http://www.statehealthfacts.org/comparebar.jsp?ind=138&cat=3> on November 8, 2010.
9. Centers for Disease Control and Prevention. 2012. Diagnoses of HIV Infection and AIDS in the United States and Dependent Areas, 2010. HIV Surveillance Report, Volume 22. Accessed on July 5, 2013 at <http://www.cdc.gov/hiv/surveillance/resources/reports/2010report/index.htm>.
10. CDC. Summary Health Statistics for the U.S. Population, National Health Interview Survey, 2012. Accessed from http://www.cdc.gov/nchs/data/series/sr_10/sr10_248.pdf on October 26, 2014.
11. Sexually Transmitted Disease Surveillance 2012. Atlanta: U.S. Department of Health and Human Services; 2013. Accessed March 11, 2014.
12. Gay Lesbian Straight Education Network (GLSEN). The 2011 National School Climate Survey. Accessed from <http://www.glsen.org/cgi-bin/iowa/all/library/record/2624.html?state=research&type=research> on August 16, 2014.
13. Task Force. Lesbian, Gay, Bisexual, and Transgender Youth: An Epidemic of Homelessness. Washington DC: National Gay and Lesbian Task Force, 2006.
14. U.S. House of Representatives, Committee on Government Reform. The Content of Federally Funded Abstinence-Only Education Programs, Prepared for Rep. Henry A. Waxman. Washington, DC: Author, 2004.
15. Brooks RA et al. "Preventing HIV among Latino and African American Gay and Bisexual Men in a Context of HIV-Related Stigma, Discrimination, and Homophobia: Perspectives of Providers." *AIDS Patient Care STDS*. 2005 November; 19(11): 737-744.
16. Fortenberry JD et al. "Relationships of stigma and shame to gonorrhea and HIV screening." *Am J Public Health*. 2002 Mar;92(3):378-81.
17. Huebner DM et al. The Impact of Internalized Homophobia on HIV. *American Journal of Community Psychology*, June 2002; 30(3) 327-348. Preventive Interventions *American Journal of Community Psychology*, Vol. 30, No. 3, June 2002.
18. Centers for Disease Control and Prevention (CDC). "National HIV Behavioral Surveillance: HIV Risk and Testing Behaviors Among Young MSM." Accessed from <http://www.cdc.gov/hiv/topics/msm/ymsm.htm> on 8/6/2014.
19. Hurt CB, Matthews DD, Calabria BM, et al. Sex with older partner is associated with primary HIV infection among men who have sex with men in North Carolina. *J Acquir Immune Defic Syndr*. 2010;54:185-190.
20. CDC. "HIV Risk, Prevention, & Testing Behaviors among MSM National HIV Behavioral Surveillance System" *MMWR*, October 1, 2014.
21. MacKellar DA et al. "A Plausible Causal Model of HAART-Efficacy Beliefs, HIV/AIDS Complacency, and HIV-Acquisition Risk Behavior Among Young Men Who Have Sex with Men." *AIDS Behav*, Published Online September 23, 2010.
22. Kelly, J. A., Murphy, D. A., Sikkema, K. J., McAuliffe, R. L., Roffman, R. A., Solomon, L. J., Winett, R. A., Kalichman, S. C. (1997). Randomised, controlled, community-level HIV-prevention intervention for sexual-risk behaviour among homosexual men in US cities. *Community HIV Prevention Research Collaborative*. *Lancet*, 350(9090), 1500-1505.
23. Kegeles SM, Hays RB, Coates TJ. The Mpowerment Project: a community-level HIV prevention intervention for young gay and bisexual men. *Am J Public Health* 1996; 86:1129-1136.
24. Remafedi G. Cognitive and behavioral adaptations to HIV/AIDS among gay and bisexual adolescents. *J Adolesc Health* 1994; 15:142-148.
25. Wilton, L., Herbst, J. H., Coury-Doniger, P., Painter, T. M., English, G., Alvarez, M. E., et al. (2009). Efficacy of an HIV/STI prevention intervention for black men who have sex with men: Findings from the Many Men, Many Voices (3MV) Project. *AIDS and Behavior*, 13, 532-544.
26. Beeker C, Kraft JM, Peterson JL, et al. Influences on sexual risk behavior in young African American men who have sex with men. *J Gay & Lesbian Medical Assoc* 1998; 2:59-67.
27. Choi KH, Yap GA, Kumekawa E. HIV prevention among Asian and Pacific Islander men who have sex with men: a critical review of theoretical models and directions for future research. *AIDS Educ Prev* 1998; 10 (Suppl A):19-30.
28. Gavin, Loretta E. et al. A Review of Positive Youth Development Programs That Promote Adolescent Sexual And Reproductive Health. *Journal of Adolescent Health*. 46 (2010) S75.
29. Kirby D. Emerging Answers: Research Findings on Programs to Reduce Teen Pregnancy. Washington, DC: National Campaign to Prevent Teen Pregnancy, 2007.
30. Stevens J. Peer Education: Promoting Healthy Behaviors. [The Facts] Washington, DC: Advocates for Youth, 1997.
31. Kurth AE et al. Combination HIV Prevention: Significance, Challenges, and Opportunities. *Current HIV/AIDS Reports*, published online October 13, 2010.
32. Pace, J. E., Siberry, G. K., Hazra, R., & Kapogiannis, B. G. (2013). Preexposure prophylaxis for adolescents and young adults at risk for HIV infection: is an ounce of prev
33. Isbell MT. HIV Prevention for Gay Men and Men who have Sex with Men: Development of a Comprehensive Policy Agenda. *amfAR and Trust for America's Health*, 2010.
34. DiClemente RJ, Salazar LF, Crosby RA. A Review of STD/HIV Preventive Interventions for Adolescents: Sustaining Effects Using an Ecological Approach. *Journal of Pediatric Psychology* 2007; 32(8): 888-906.

MISSION

Advocates for Youth partners with youth leaders, adult allies, and youth-serving organizations to advocate for policies and champion programs that recognize young people's rights to: honest sexual health information; accessible, confidential, and affordable sexual health services; and the resources and opportunities necessary to create sexual health equity for **all** youth.

OUR VISION: THE 3RS

Advocates' vision is informed by its core values of ***Rights.Respect.Responsibility***.

Advocates believe that:

RIGHTS Youth have the inalienable ***right*** to honest sexual health information; confidential, consensual sexual health services; and equitable opportunities to reach their full potential.

RESPECT Youth deserve ***respect***. Valuing young people means authentically involving them in the design, implementation, and evaluation of programs and policies that affect their health and well-being.

RESPONSIBILITY Society has the ***responsibility*** to provide young people with all of the tools they need to safeguard their sexual health, and young people have the responsibility to protect themselves.

SOME RELATED PUBLICATIONS FROM ADVOCATES FOR YOUTH

The Facts: Young People and HIV in the United States

The Facts: Young Men Who Have Sex With Men: At Risk for HIV and STIs

The Facts: LGBTQ Youth

See the complete library of publications at www.advocatesforyouth.org/publications